Labcorp Oklahoma, Inc. Test Directory

Calcitonin

Order Name: CALCITONI
Test Number: 3600550
Revision Date: 12/12/2022

| 49482). Freeze immediately and maintain frozen until tested. To avoid delays in turnaround time when requesting multiple tests on frozen sampl PLEASE SUBMIT SEPARATE FROZEN SPECIMENS FOR EACH TEST REQUESTED. Special Instructions: State on the request form if calcium infusion or pentagastrin injection tests are part of the patient preparation. Values obtawith different assay methods should not be used interchangeably in serial testing. It is recommended that only one assay method be used consistently to monitor each patient's course of therapy. This procedure does not provide serial monitoring; it is intended for one-time use only. | TEST NAME | | | METHODOLOGY | LOINC CODE | |
|---|---------------|--|---------------|---|-----------------------|--|
| Specimen Volume (min) Specimen Type Specimen Container Transport Environment Preferred 1 mL (0.4) Serum Clot Activator SST Frozen Instructions Notes: 0.4 mL (Note: This volume Does NOT allow for repeat testing.) Specimen Type: Red-top tube or gel-barrier tube Container Detail: LC PP Specimen Storage: Freeze Specimen Collection: Separate serum from cells. Transfer the serum into a LabCorp PP transpak frozen purple tube with screw cap (LabCorp 49482). Freeze immediately and maintain frozen until tested. To avoid delays in turnaround time when requesting multiple tests on frozen sample PLEASE SUBMIT SEPARATE FROZEN SPECIMENS FOR EACH TEST REQUESTED. Special Instructions: State on the request form if calcium infusion or pentagastrin injection tests are part of the patient preparation. Values obtain with different assay methods should not be used interchangeably in serial testing. It is recommended that only one assay method be used consistently to monitor each patient's course of therapy. This procedure does not provide serial monitoring; it is intended for one-time use only. | Calcitonin | | | Quantitative Chemiluminescent Immunoass | 1992-7 | |
| Preferred 1 mL (0.4) Serum Clot Activator SST Frozen Notes: 0.4 mL (Note: This volume Does NOT allow for repeat testing.) Specimen Type: Red-top tube or gel-barrier tube Container Detail: LC PP Specimen Storage: Freeze Specimen Collection: Separate serum from cells. Transfer the serum into a LabCorp PP transpak frozen purple tube with screw cap (LabCorp 49482). Freeze immediately and maintain frozen until tested. To avoid delays in turnaround time when requesting multiple tests on frozen sampled PLEASE SUBMIT SEPARATE FROZEN SPECIMENS FOR EACH TEST REQUESTED. Special Instructions: State on the request form if calcium infusion or pentagastrin injection tests are part of the patient preparation. Values obtain with different assay methods should not be used interchangeably in serial testing. It is recommended that only one assay method be used consistently to monitor each patient's course of therapy. This procedure does not provide serial monitoring; it is intended for one-time use only. | SPECIMEN REQU | IREMENTS | | | | |
| Instructions Notes: 0.4 mL (Note: This volume Does NOT allow for repeat testing.) Specimen Type: Red-top tube or gel-barrier tube Container Detail: LC PP Specimen Storage: Freeze Specimen Collection: Separate serum from cells. Transfer the serum into a LabCorp PP transpak frozen purple tube with screw cap (LabCorp 49482). Freeze immediately and maintain frozen until tested. To avoid delays in turnaround time when requesting multiple tests on frozen sample PLEASE SUBMIT SEPARATE FROZEN SPECIMENS FOR EACH TEST REQUESTED. Special Instructions: State on the request form if calcium infusion or pentagastrin injection tests are part of the patient preparation. Values obtain with different assay methods should not be used interchangeably in serial testing. It is recommended that only one assay method be used consistently to monitor each patient's course of therapy. This procedure does not provide serial monitoring; it is intended for one-time use only. | Specimen | Specimen Volume (min) | Specimen Type | Specimen Container | Transport Environment | |
| Specimen Type: Red-top tube or gel-barrier tube Container Detail: LC PP Specimen Storage: Freeze Specimen Collection: Separate serum from cells. Transfer the serum into a LabCorp PP transpak frozen purple tube with screw cap (LabCorp 49482). Freeze immediately and maintain frozen until tested. To avoid delays in turnaround time when requesting multiple tests on frozen sampl PLEASE SUBMIT SEPARATE FROZEN SPECIMENS FOR EACH TEST REQUESTED. Special Instructions: State on the request form if calcium infusion or pentagastrin injection tests are part of the patient preparation. Values obtawith different assay methods should not be used interchangeably in serial testing. It is recommended that only one assay method be used consistently to monitor each patient's course of therapy. This procedure does not provide serial monitoring; it is intended for one-time use only. | Preferred | 1 mL (0.4) | Serum | Clot Activator SST | Frozen | |
| Container Detail: LC PP Specimen Storage: Freeze Specimen Collection: Separate serum from cells. Transfer the serum into a LabCorp PP transpak frozen purple tube with screw cap (LabCorp 49482). Freeze immediately and maintain frozen until tested. To avoid delays in turnaround time when requesting multiple tests on frozen sampl PLEASE SUBMIT SEPARATE FROZEN SPECIMENS FOR EACH TEST REQUESTED. Special Instructions: State on the request form if calcium infusion or pentagastrin injection tests are part of the patient preparation. Values obtain with different assay methods should not be used interchangeably in serial testing. It is recommended that only one assay method be used consistently to monitor each patient's course of therapy. This procedure does not provide serial monitoring; it is intended for one-time use only. | Instructions | Notes: 0.4 mL (Note: This volume Does NOT allow for repeat testing.) | | | | |
| Specimen Storage: Freeze Specimen Collection: Separate serum from cells. Transfer the serum into a LabCorp PP transpak frozen purple tube with screw cap (LabCorp 49482). Freeze immediately and maintain frozen until tested. To avoid delays in turnaround time when requesting multiple tests on frozen sampl PLEASE SUBMIT SEPARATE FROZEN SPECIMENS FOR EACH TEST REQUESTED. Special Instructions: State on the request form if calcium infusion or pentagastrin injection tests are part of the patient preparation. Values obtawith different assay methods should not be used interchangeably in serial testing. It is recommended that only one assay method be used consistently to monitor each patient's course of therapy. This procedure does not provide serial monitoring; it is intended for one-time use only. | | Specimen Type: Red-top tube or gel-barrier tube | | | | |
| Specimen Collection: Separate serum from cells. Transfer the serum into a LabCorp PP transpak frozen purple tube with screw cap (LabCorp 49482). Freeze immediately and maintain frozen until tested. To avoid delays in turnaround time when requesting multiple tests on frozen sampl PLEASE SUBMIT SEPARATE FROZEN SPECIMENS FOR EACH TEST REQUESTED. Special Instructions: State on the request form if calcium infusion or pentagastrin injection tests are part of the patient preparation. Values obtain with different assay methods should not be used interchangeably in serial testing. It is recommended that only one assay method be used consistently to monitor each patient's course of therapy. This procedure does not provide serial monitoring; it is intended for one-time use only. | | Container Detail: LC PP | | | | |
| 49482). Freeze immediately and maintain frozen until tested. To avoid delays in turnaround time when requesting multiple tests on frozen sampl PLEASE SUBMIT SEPARATE FROZEN SPECIMENS FOR EACH TEST REQUESTED. Special Instructions: State on the request form if calcium infusion or pentagastrin injection tests are part of the patient preparation. Values obtawith different assay methods should not be used interchangeably in serial testing. It is recommended that only one assay method be used consistently to monitor each patient's course of therapy. This procedure does not provide serial monitoring; it is intended for one-time use only. | | Specimen Storage: Freeze | | | | |
| PLEASE SUBMIT SEPARATE FROZEN SPECIMENS FOR EACH TEST REQUESTED. Special Instructions: State on the request form if calcium infusion or pentagastrin injection tests are part of the patient preparation. Values obtawith different assay methods should not be used interchangeably in serial testing. It is recommended that only one assay method be used consistently to monitor each patient's course of therapy. This procedure does not provide serial monitoring; it is intended for one-time use only. | | Specimen Collection: Separate serum from cells. Transfer the serum into a LabCorp PP transpak frozen purple tube with screw cap (LabCorp No | | | | |
| Special Instructions: State on the request form if calcium infusion or pentagastrin injection tests are part of the patient preparation. Values obtation with different assay methods should not be used interchangeably in serial testing. It is recommended that only one assay method be used consistently to monitor each patient's course of therapy. This procedure does not provide serial monitoring; it is intended for one-time use only. | | 49482). Freeze immediately and maintain frozen until tested. To avoid delays in turnaround time when requesting multiple tests on frozen samples, | | | | |
| with different assay methods should not be used interchangeably in serial testing. It is recommended that only one assay method be used consistently to monitor each patient's course of therapy. This procedure does not provide serial monitoring; it is intended for one-time use only. | | PLEASE SUBMIT SEPARATE FROZEN SPECIMENS FOR EACH TEST REQUESTED. | | | | |
| consistently to monitor each patient's course of therapy. This procedure does not provide serial monitoring; it is intended for one-time use only. | | Special Instructions: State on the request form if calcium infusion or pentagastrin injection tests are part of the patient preparation. Values obtained | | | | |
| | | with different assay methods should not be used interchangeably in serial testing. It is recommended that only one assay method be used | | | | |
| 0 1 0 177 1 1 1 1 1 1 1 1 5 1 1 1 1 1 5 1 1 1 1 | | consistently to monitor each patient's course of therapy. This procedure does not provide serial monitoring; it is intended for one-time use only. | | | | |
| Specimen Stability: Ambient: Unstable, Retrigerated : Unstable, Frozen: 90 days | | Specimen Stability: Ambient: Unstable, Refrigerated : Unstable, Frozen: 90 days | | | | |

| GENERAL INFORMATION | | |
|---------------------------------|---------------------------|--|
| Expected TAT | 3 - 5 days | |
| Performing Labcorp Test Code | 004895 | |
| Notes | Labcorp Test Code: 004895 | |
| CPT Code(s) | 82308 | |
| Lab Section | Reference Lab | |

Service provided by Labcorp Oklahoma, Inc. All Rights Reserved. © 2003 - 2025