

Cystatin C

Order Name: **CYSTATIN C**

Test Number: 3623775

Revision Date: 10/29/2025

TEST NAME	METHODOLOGY	LOINC CODE
Cystatin C	Turbidometric	33863-2

SPECIMEN REQUIREMENTS				
Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Preferred	1 mL (0.5 mL)	Serum	Clot Activator SST	Refrigerated
Alternate 1	1 mL (0.5 mL)	Plasma	Sodium Heparin (Green Top / No-Gel)	Refrigerated
Instructions	Dietary Instructions: Overnight fasting is preferred. Collect blood in SST Clot tube or Sodium Heparin Green-top tube. Allow Serum to clot for 15 minutes. Centrifuge aliquot Serum or Plasma for 10min and separate wihtin 2 hours and keep refrigerated. Specimen Stability: Room temperature: 24 Hours, Refrigerated: 7 Days, Frozen: 90 Days.			

GENERAL INFORMATION				
Testing Schedule	Mon-Fri			
Expected TAT	3-4 Days			
Clinical Use	Cystatin C is a highly sensitive and specific marker of glomerular filtration rate (renal function). Cystatin C is independent of muscle mass, age, and body mass index. Cystatin C is also used to assess renal allograft function.			
Notes	This test will order and result a Cystatin C Estimated GFR as shown below.			
	ORDER CODE	REFLEX NAME	RESULT CODE	RESULT LONG NAME
	3623778	.Cystatin C Estimated GFR	3623888	Cystatin C eGFR
CPT Code(s)	82610			
Lab Section	Immunology - Serology			