

Striated Muscle Antibody Screen with Titer

Order Name: **STRIAT AB**
Test Number: 3805400
Revision Date: 03/10/2026

TEST NAME	METHODOLOGY	LOINC CODE
Striated Muscle Antibody Screen	Semi-Quantitative Indirect Fluorescent Antibody	5372-8

SPECIMEN REQUIREMENTS

Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Preferred	1 mL (0.5 mL)	Serum	Clot Activator SST	Refrigerated
Alternate 1	1 mL (0.5 mL)	Serum	Clot Activator (Red Top, No-Gel)	Refrigerated

Instructions

Minimum Volume: 0.5 mL (Note: This volume does not allow for repeat testing.)
Specimen Type: Gel-barrier tube or Red-top tube
Specimen Storage: Refrigerated
Specimen Collection: Separate serum from cells ASAP or within 2 hours of collection
Specimen Stability: Ambient: 14 days, Refrigerated : 14 days, Frozen: 14 days (Freeze/thaw cycles Stable x3)
Causes for Rejection: Hemolyzed or lipemic serum

If being collected for Lambert-Eaton Syndrome Panel, Please collect Serum from Red No-Gel Clot Tube.

GENERAL INFORMATION

Expected TAT	2 - 4 days
Clinical Use	This test is recommended for differential evaluation of neuromuscular junction diseases including myasthenia gravis (MG). This test will report a Negative Screen and if Positive will report out the Titer. Reference Interval Negative: <1:100
Performing Labcorp Test Code	160184
CPT Code(s)	86255
Lab Section	Reference Lab