Parathyroid Hormone (PTH), Intraoperative, Baseline

NRLS-Core Chemistry

Lab Section

Order Name: **PTHIOBL**Revision Date: 04/20/2015

TEST NAME			METHODOLOGY	LOINC CODE
Parathyroid Hormone	(PTH), Intraoperative, Baseline			
SPECIMEN REQUIRE	MENTS			
Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Preferred	4 mL (2 mL)	Whole Blood	Lavender Top K2 EDTA	Refrigerated
Alternate 1	2 mL (1 mL)	Plasma	Lavender Top K2 EDTA	Frozen
Alternate 2	2 mL (1 mL)	Serum	Serum Gel Tube	Frozen
Instructions	Draw blood in a lavender-top (K2 EDTA) tube(s), avoid hemolysis. Alternative container: Plasma collected in EDTA tube, frozen; Serum collected in gold gel or plain red tube, frozen. Note: Label specimen with patient name (first and last) or other unique identifier, patient hospital identification number or Social Security number, date and time of draw, collector initials, and test(s) being ordered. Note: Intraoperative PTH should be drawn in Ambulatory Care Unit or Surgery Holding.			
Reference Range	See Interpretive Report			
Methodology	Immunoenzymatic			
GENERAL INFORMA	TION			
Testing Schedule	Sunday through Saturday			
Expected TAT	3 Hours			
Stat TAT	45 Minutes			
CPT Code(s)	83970			